

The following article is ideal because it fully supports our theory for elements as the foundation for cell regulation in conjunction with amino acids.

The only missing piece in this work is the third “electrolyte” of calcium - zinc and the details for titin; the “gene” that encompasses twelve cytokines that regulate mitochondria for cardiomyocytes and others that utilize gasotransmitters to form cell surface signaling molecules that enter cells through endocytosis.

The following are provided for discussion purposes:

Cell Surface Formation

<http://www.mcfip.net/upload/Cell%20Surface%20Signaling%20Molecule%20Formation%207-2017.pdf>

Endocytosis

<http://www.mcfip.net/upload/Endocytosis%20Modeling%204-30-17.pdf>

Summary

Biomedical research has evolved to ignore the role of elements as the foundation for signaling within and between cells. As a result, the primary causes of chronic diseases have not been identified.

The following is provided for discussion purposes.

[http://www.mcfip.net/upload/Crystallography%20Overview%20x%20\(1\).pdf](http://www.mcfip.net/upload/Crystallography%20Overview%20x%20(1).pdf)

Using the fundamentals of physical science and the interactions between elements our epigenetic modeling methods identify primary causes of chronic diseases. This assertion can be verified by using the links provided in the tabs of the MCFIP website: [www.MCFIP.net](http://www.MCFIP.net).

# Force field analysis provides clues to protein-ion interaction

October 17, 2017

Muscle twitching, lactose digestion, blood movement—what could possibly connect these body functions? You may be surprised to learn that all these processes and many more are driven by metal ions.

Sodium (Na<sup>+</sup>), potassium (K<sup>+</sup>), calcium (Ca<sup>2+</sup>) and magnesium (Mg<sup>2+</sup>) might be household names, but they are also critical to the function of human cells. The combination of these ions with bodily proteins creates complexes that are imperative to our continued existence.

The importance of proteins and metal ion interactions is well understood, but the mechanistic interactions between the two are still far from a complete picture. Zhifeng Jine, Rui Qi, Chengwen Liu and Pengyu Ren, professors in the biomedical engineering department at the University of Texas at Austin, are working to quantitatively describe protein-ion interactions using what is called an atomic multipole optimized energetics for biomolecular applications (AMOEBA) force field. They describe their work in this week's the *Journal of Chemical Physics*.

AMOEBA technology was developed by Ren and Jay Ponder, from the University of Texas at Austin and Washington University in St. Louis. It uses a polarizable atomic multipole model to calculate the potential energy of a system. The AMOEBA model aims to address known inadequacies of current force field technology to accurately model ion and protein interactions.

"The lack of detailed understanding is mainly due to the lack of accurate and yet computationally efficient models to treat metal ions," Ren said. "We aim to apply high-level quantum mechanical methods and advanced force field simulations to understand the nature of the interactions between metal ions and proteins."

AMOEBA improves on classically based biomolecular simulations, whose models represent the intermolecular interaction using the sum of two forces, according to Ren: the van der Waals interaction and the Coulomb interaction between point charges. "This representation is simple enough to be computationally tractable, and it has been widely used over the last few decades," he said. "However, metal ions can lead to strong polarization and charge transfer effects, which are missing in these classical models. We believe these effects play [a] critical role in the specific ion-protein interaction."

Ren and his colleagues looked specifically at the interaction of  $Mg^{2+}/Ca^{2+}$  and amino acids. Magnesium and calcium are some of the most abundant ions in metalloproteins, proteins with a metal ion cofactor. Both are selectively bound to specific amino acids, making them interesting target molecules. The many-body response, the mutual induction between ions and their surrounding residues within the protein binding pockets, also impacts the binding affinity of these ion and can be captured by the AMOEBA model.

"The difference between the many-body response to  $Ca^{2+}$  vs  $Mg^{2+}$  in the binding pocket is substantial," said Ren. "It was known that polarization and charge-transfer are important in protein-ion complexes, but for the competitive binding, many have suspected that these effects might cancel out."

The AMOEBA model and the advancements of force fields, including applications of these results, Ren pointed out, are relevant in many diseases including cancer and neurodegenerative disorders. Knowledge of protein-ion interactions can provide foundational understanding for the advancement in related medical researches.

**Explore further: Travelling-wave ion mobility mass spectrometry elucidates structures of gold fingers**

**More information:** "Study of interactions between metal ions and protein model compounds by energy decomposition analyses and the AMOEBA force field," *Journal of Chemical Physics* (2017). DOI: [10.1063/1.4985921](https://doi.org/10.1063/1.4985921)

**Journal reference:** *Journal of Chemical Physics*

Read more at: <https://phys.org/news/2017-10-field-analysis-clues-protein-ion-interaction.html#jCp>